

Improvements in Implant Dentistry over the Last Decade: Comparison of Survival and Complication Rates in Older and Newer Publications

Bjarni E. Pjetursson, DDS, Dr Med Dent, MAS (Perio)¹/Asgeir G. Asgeirsson, DDS^{1,3}/
Marcel Zwahlen, MSc, PhD²/Irena Sailer, Dr Med Dent³

Purpose: The objective of this systematic review was to assess and compare the survival and complication rates of implant-supported prostheses reported in studies published in the year 2000 and before, to those reported in studies published after the year 2000. **Materials and Methods:** Three electronic searches complemented by manual searching were conducted to identify 139 prospective and retrospective studies on implant-supported prostheses. The included studies were divided in two groups: a group of 31 older studies published in the year 2000 or before, and a group of 108 newer studies published after the year 2000. Survival and complication rates were calculated using Poisson regression models, and multivariable robust Poisson regression was used to formally compare the outcomes of older and newer studies. **Results:** The 5-year survival rate of implant-supported prostheses was significantly increased in newer studies compared with older studies. The overall survival rate increased from 93.5% to 97.1%. The survival rate for cemented prostheses increased from 95.2% to 97.9%; for screw-retained reconstruction, from 77.6% to 96.8%; for implant-supported single crowns, from 92.6% to 97.2%; and for implant-supported fixed dental prostheses (FDPs), from 93.5% to 96.4%. The incidence of esthetic complications decreased in more recent studies compared with older ones, but the incidence of biologic complications was similar. The results for technical complications were inconsistent. There was a significant reduction in abutment or screw loosening by implant-supported FDPs. On the other hand, the total number of technical complications and the incidence of fracture of the veneering material was significantly increased in the newer studies. To explain the increased rate of complications, minor complications are probably reported in more detail in the newer publications. **Conclusions:** The results of the present systematic review demonstrated a positive learning curve in implant dentistry, represented in higher survival rates and lower complication rates reported in more recent clinical studies. The incidence of esthetic, biologic, and technical complications, however, is still high. Hence, it is important to identify these complications and their etiology to make implant treatment even more predictable in the future. INT J ORAL MAXILLOFAC IMPLANTS 2014;29(SUPPL):308–324. doi: 10.11607/jomi.2014suppl.g5.2

Key words: biologic complications, dental implants, esthetic complications, failures, fixed dental prosthesis, implant dentistry, marginal bone loss, single crowns, survival, success, systematic review, technical complications

In recent years, a number of systematic reviews on the survival and complication rates of fixed implant-

supported prostheses have been published. The reviews focused on implant-supported single crowns (SCs)^{1,2} or on multiple-unit implant-supported fixed dental prostheses (FDPs)^{3,4,5} and reported high 5- and 10-year survival rates for both types of prostheses. Due to these good results, fixed implant-supported prostheses are fully accepted as a reliable treatment option for the replacement of single or multiple missing teeth today.^{1,4}

In the daily clinical practice, however, patient (and clinician) satisfaction is not only influenced by survival rates. Survival rates in general represent prostheses that remained in clinical service for a defined follow-up period. However, those prostheses were not necessarily free of complications. Both implant-supported

¹University of Iceland, Faculty of Odontology, Reykjavik, Iceland.

²University of Berne, Department of Social and Preventive Medicine, Bern, Switzerland.

³University of Zürich, Department of Fixed and Removable Prosthodontics and Dental Material Science, Zürich, Switzerland.

Correspondence to: Dr Bjarni E. Pjetursson, Department of Reconstructive Dentistry, University of Iceland, Vatnsmyrarvegur 16, IS-101 Reykjavik, Iceland. Email: bep@hi.is

©2014 by Quintessence Publishing Co Inc.

SCs as well as implant-supported FDPs suffer from different kinds of biologic or technical complications as the reviews indicated.^{1,5} These complications lead to the need for corrective treatment, increasing the total chairside time and the treatment costs. As a consequence, a reduction of the general satisfaction with the prosthesis may occur.⁶

The etiology of biologic complications is mostly patient-based and can be multifactorial (eg, hereditary susceptibility to peri-implantitis, bad oral hygiene, excess cement).⁷ A reduction of risk therefore implies good patient compliance and intensive oral care. Biologic risk can only be minimally influenced by modification of the implants, implant surfaces, and components, according to the current literature.

Technical problems are mostly related to the materials and the design of the components.⁸ Studies have shown various types of technical problems, like prosthetic fixation screw or abutment loosening, fractures of components (eg, abutments, screws), fractures of reconstructive materials (eg, chipping of veneering ceramic), and loss of retention of cemented prostheses due to fracture of the luting cement.^{1,5} In contrast to the biologic risks, the technical outcome can be improved with technical amendments.

In order to reduce the risk for technical complications, the materials and components used for the implant-supported prostheses are, therefore, constantly being enhanced. Some improvements have already led to better outcomes. As an example, after the introduction of implant-supported SCs, very high numbers of abutment or occlusal screw loosening were reported.^{9,10} The change of screw material from titanium to gold and the use of defined screw fixation torques led to significant lowering of the incidence of screw loosening.^{9,10} Screw loosening is still one of the most frequently reported complications for implant-supported reconstruction. Therefore, further refinements are desired, and debates about the best materials and techniques for the implant-supported prostheses are continually raised.^{1,4}

The introduction of new restorative materials, such as ceramic zirconia for the abutment and framework, can on the one hand improve the outcomes (esthetics), but on the other increase technical problems. It has been shown that the veneering ceramics for zirconia-based prostheses exhibited very high rates for fracture and chipping.¹¹⁻¹³ Thus, not all further developments were really an improvement.

Very little scientific evidence is currently available to help determine whether changes in materials and implant components in the last decades have influenced survival rates of implants and implant-supported prostheses and the incidence of biologic and technical complications.

The objective of this systematic review was to assess and compare the survival and complication rates of implant-supported prostheses reported in studies published in the year 2000 and before to those reported in studies published after the year 2000.

MATERIALS AND METHODS

Focus Question

The following focus question was developed using a PICO approach:

Have the survival rate of implant-supported prostheses and the incidence of complications changed over the last decade?

Search Strategy and Study Selection

Three Medline (PubMed) searches were performed for articles published in the Dental Literature in English and German. The first one covered the time period 1990 to September 2012 utilizing both MeSH terms and free text words. The following search terms were used: "implant*", "cement*" or "screw*", "fix*" or "retain*", "single-crown", "single crowns" "FPD", "FDP", "bridge", "reconstruct*" and "suprastruct*". Moreover, the terms "long-term", "long term", "longitud*", "survival" or "failure", "complicat*", "technical" or "biological" were utilized. In addition, Cochrane Library and Embase searches were conducted applying the same search terms.¹⁰ The second search was an updated search from a previous systematic review¹ covering the time interval through the end of August 2011. The following MeSH terms were selected for the search: "dental implants" AND ("crowns" OR "survival").² The third search was performed for studies published between May 1, 2004, and August 31, 2011,⁵ using the following MeSH search terms: "dental implants" AND ("denture, partial, fixed" OR survival). Additionally, the studies from the predecessor systematic review were included, encompassing publications from 1966 through the end of April 2004.¹⁴ All three searches were complemented by manual searches of the bibliographies of all full-text articles and related reviews, selected from the electronic search. Furthermore, manual searching was applied to relevant journals in the field of interest (Table 1).

Inclusion Criteria

This systematic review was based on randomized controlled clinical trials (RCTs), controlled clinical trials (CCTs), prospective cohort studies, prospective case series, and retrospective studies. The additional inclusion criteria for study selection were:

Table 1 Search Strategy

Focus question Have the survival rate of implant-supported reconstructions and the incidence of complications changed over the last decade?

Search strategy

Population	Partially edentulous patients with single-implant FDPs or multi-unit partial/full FDPs
Intervention or exposure	1 year of clinical follow-up, and after 5 and 10 y of follow-up
Comparison	Different decades/timepoints of intervention and/or publication
Outcome	Survival and complication rates over time
Search combination	Sailer et al ¹⁰ : used "implant*", "cement*" or "screw*", "fix*" or "retain*", "single-crown", "single crowns" "FPD", "FDP", "bridge", "reconstruct*" and "suprastruct*". Moreover, the terms "long-term", "long term", "longitud*", "survival" or "failure", "complicat*", "technical" or "biological" Jung et al ² : used "dental implants" and ("crowns" OR "survival"). Pjetursson et al ⁵ : used "dental implants" AND ("denture, partial, fixed" OR survival). All these search terms were MeSH terms.

Database search

Electronic	Medline (PubMed), Cochrane Library, and Embase
Journals	<i>Clinical Oral Implants Research, International Journal of Oral and Maxillofacial Implants, International Journal of Oral Surgery, Journal of Prosthetic Dentistry, International Journal of Periodontics and Restorative Dentistry, International Journal of Prosthodontics, Clin Implant Dent Relat Res, Journal of Clinical Periodontology, Journal of Oral Rehabilitation, British Dental Journal, Journal of Prosthodontics, Implant Dentistry, Journal of Periodontology, International Journal of Epidemiology, European Journal of Oral Sciences, Australian Dental Journal, Clinical Oral Investigations, Dental Materials, Journal of the Canadian Dental Association, Quintessence International, Journal of Oral Implantology, Clinical Implant Dentistry and Related Research</i>

Selection criteria

Inclusion criteria	This systematic review was based on randomized controlled clinical trials (RCTs), controlled clinical trials (CCTs), prospective cohort studies, prospective case series, and retrospective studies. The additional inclusion criteria for study selection were: Human trials with minimum of 10 subjects. Studies reported in English and German language and published in dental journals. Patients were examined clinically at the follow-up visit. Publications based on patient records only, on questionnaires or interviews were excluded. Studies reported details on the characteristics and outcome of the suprastructures. For the short-term data, the studies had to have a mean time of functional loading of at least 1 year. For the longitudinal data, the studies had to have a mean follow-up time of 5 years or more.
Exclusion criteria	The main reasons for exclusion were lack of detailed information on the reconstruction design and no detailed information on the outcome of the reconstruction at the follow-up visit, or mean observation period not fulfilling the inclusion criteria and studies with less than 10 subjects. Furthermore, publications based on questionnaires or interviews without clinical examinations, multiple publications on the same patient cohorts, and case descriptions of failures without relevant information on the entire patient cohort were excluded.

- Human trials with a minimum of 10 subjects.
- Studies reported in English and German language and published in dental journals.
- Patients examined clinically at the follow-up visit. Publications based on patient records only, on questionnaires or interviews were excluded.
- Studies reporting details on the characteristics and outcome of the suprastructures.
- For short-term data, the studies had to have a mean time of functional loading of at least 1 year.
- For longitudinal data, the studies had to have a mean follow-up time of 5 years or more.

Selection of Studies

Titles and abstracts of the searches were always screened by at least two independent reviewers for possible inclusion in the reviews. The full text of all studies of possible relevance was then obtained for independent assessment by the reviewers. Any disagreement was resolved by discussion.

The first search extending through August 2011 identified 59 full-text articles that gave information on the clinical performance of cemented and screw-retained implant-supported prostheses with a functional loading of at least 1 year.¹⁰ The extended search

up to September 2012 identified two additional publications fulfilling the inclusion criteria. In the second search, the original search extending until August 2006 identified 24 studies reporting on implant-supported single crowns (SCs) with a mean follow-up time of 5 years or more.¹ The extended search, through August 2011, added 22 new publications to the included studies.² In the third search, identifying implant-supported fixed dental prosthesis (FDPs) with a mean follow-up time of at least 5 years, the original search extending until May 2004 identified 21 studies.¹⁴ The extended search through August 2011 identified an additional 11 studies.⁵

The results of the present systematic review are based on a total of 139 included studies.

Excluded Studies

The main reasons for exclusion were no detailed information on the prostheses design and no detailed information on the outcome of the prosthesis at the follow-up visit, mean observation period not fulfilling the inclusion criteria, and studies with less than 10 subjects. Furthermore, publications based on questionnaires or interviews without clinical examinations, multiple publications on the same patient cohorts, and case descriptions of failures without relevant information on the entire patient cohort were excluded.

Data Extraction

From the included studies, information on failures of the supporting implants and the prostheses was extracted. Information on esthetic, biologic, and technical complications was also retrieved. Biologic complications were characterized by a biological process affecting the supporting tissues. Soft tissue complications and peri-implantitis characterized by a substantial (> 2 mm) marginal bone loss were included in this category.

Technical complications were characterized by mechanical damage of implants, abutments, and/or the suprastructures. Among these, fractures of implants, screws, or abutments; fractures of the luting cement (loss of retention); fractures or deformations of the framework or veneers; and screw or abutment loosening were included. From the included studies, the number of events for all of these categories was abstracted and the corresponding total exposure time of the implants, abutments, and prostheses was calculated.

Statistical Analysis

By definition, failure and complication rates are calculated by dividing the number of events (failures or complications) in the numerator by the total exposure time (implant, abutment, or reconstruction time) in the denominator.

The numerator could usually be extracted directly from the publication. The total exposure time was calculated by taking the sum of:

- Exposure time of implants, abutments, or prostheses that could be followed for the entire observation time.
- Exposure time up to a failure of implants, abutments, or prostheses that were lost due to failure during the observation time.
- Exposure time up to the end of observation time for implants, abutments, or prostheses that did not complete the observation period due to reasons such as death, change of address, refusal to participate, non-response, chronic illness, missed appointments, and work commitments.

For each study, event rates for implants, abutments, or prostheses were calculated by dividing the total number of events by the total implant, abutment, or prosthesis exposure time in years. For additional analysis, the total number of events was considered to be Poisson distributed for a given sum of abutment exposure years, and robust Poisson regression with a logarithmic link-function and total exposure time per study as an offset variable was used.¹⁵ Robust Poisson regression allowed calculation of standard errors and 95% confidence intervals, which incorporated heterogeneity among studies.

Five-year survival proportions were calculated via the relationship between event rate and survival function S , $S(T) = \exp(-T * \text{event rate})$, by assuming constant event rates.¹⁶ The 95% confidence intervals for the survival proportions were calculated by using the 95% confidence limits of the event rates. Multivariable robust Poisson regression was used to formally compare publication years and to assess other study characteristics. All analyses were performed using Stata version 12.

RESULTS

Study Characteristics

The 139 included studies were divided into three categories. The first group was a group of 61 studies reporting on the clinical performance of cemented and screw-retained implant-supported prostheses of different types. This group included 37 studies reporting on implant-supported single crowns (SCs), 16 studies that reported on implant-supported partial fixed dental prostheses (FDPs), and 18 studies reporting on implant-supported fixed complete dentures (FCDs), with various mean follow-up periods ranging from 1 to 10 years.¹⁰ The result for this group is referred to as the overall results. The second group was a group of

Table 2 Comparison of the Implant Survival Rate in Articles Published Before and After 2000

	Published before 2000		Published after 2000		P value
	Annual failure rate (95% CI)	5-year survival rate (95% CI)	Annual failure rate (95% CI)	5-year survival (95% CI)	
Overall results	0.29% (0.15–0.57)	98.6% (97.2–99.3)	0.39% (0.25–0.63)	98.1% (96.9–98.8)	.466
Implant-supported SCs	0.60% (0.39–0.90)	97.1% (95.6–98.1)	0.56% (0.40–0.78)	97.2% (96.2–98.0)	.815
Implant-supported FDPs	1.28% (1.05–1.56)	93.8% (92.5–94.9)	0.81% (0.57–1.14)	96.1% (94.4–97.2)	.021

Table 3 Comparison of the Prosthetic Survival Rate in Articles Published Through and After 2000

	Published through 2000		Published after 2000		P value
	Annual failure rate (95% CI)	5-year survival rate (95% CI)	Annual failure rate (95% CI)	5-year survival rate (95% CI)	
Overall results	1.34% (0.65–2.77)	93.5% (87.1–96.8)	0.59% (0.38–0.91)	97.1% (95.6–98.1)	.050
Cemented reconstructions	0.99% (0.58–1.69)	95.2% (91.9–97.2)	0.42% (0.23–0.75)	97.9% (96.3–98.8)	.030
Screw-retained reconstructions	5.07% (1.17–22.07)	77.6% (33.2–94.3)	0.65% (0.39–1.10)	96.8% (94.6–98.1)	.004
Implant-supported FCDs	NR	NR	0.86% (0.45–1.66)	95.8% (92.0–97.8)	NA
Implant-supported SCs	1.54% (1.01–2.34)	92.6% (88.9–95.1)	0.58% (0.35–0.95)	97.2% (95.3–98.3)	.002
Implant-supported FDPs	1.34% (0.69–2.62)	93.5% (87.7–96.6)	0.73% (0.55–0.97)	96.4% (95.3–97.3)	.087

NR = not reported; NA = not applicable.

46 studies reporting on 3,199 implant-supported SCs with a mean follow-up time of at least 5 years.² The last group was a group of 32 studies reporting on 1,881 implant-supported FDPs with a mean follow-up period of at least 5 years.⁵

The year of publication for the 139 studies (see reference list) included in this systematic review ranged from 1994 to 2012.^{2–14,17–141} Thirty-one publications were classified as older studies published in the year 2000 or before and 108 were classified as newer studies published after the year 2000. From the 61 studies reporting on overall results, the year of publication ranged from 1995 to 2012. Twelve of the studies were classified as older studies and 49 as newer studies published in the present millennium. Out of the 46 studies reporting on implant-supported SCs, the publication year ranged from 1995 to 2012. Eight of the studies were considered older studies and 38 newer studies published after the year 2000. For the 32 studies reporting on implant-supported FDPs, the year of publication ranged from 1995 to 2012. Eleven of the included studies were considered older studies, and 21 were considered newer studies published after the year 2000.

Survival

Survival was defined as the implants or prostheses remaining in situ with or without modification over the observation period.

Implant survival. The annual implant failure rates in the older publications ranged from 0.29% to 1.28%,

translating into 5-year survival rates of 93.8% to 98.6%. The annual failure rates in the newer publications ranged from 0.39% to 0.81%, translating into a 5-year survival rate of 96.1% to 98.1% (Table 2). Comparing the survival rates in the older publications with the survival rates in the newer publications, there was only a minor difference ($P = .466, .815$) of the overall results and for the implant-supported SCs. However, for the implant-supported FDPs there were significantly ($P = .021$) less implant failures in the newer studies and the 5-year implant survival rate increased from 93.8% in the older studies to 96.1% (Table 2).

Survival of Prostheses. The annual failure rate of prostheses in the older publications ranged from 0.99% to 5.07% (Figs 1 to 6), translating into a 5-year survival rate of 77.4% to 95.2%. The highest 5-year survival rate in the older studies was seen for cemented prostheses, and the lowest survival rate was reported for screw-retained prostheses (Table 3). The range in annual failure rates of different types of prostheses was significantly reduced in the newer publications. They ranged from 0.42% to 0.86%, translating into a 5-year survival rate of 95.8% to 97.9% (Table 3, Figs 1 to 6). Formally comparing the survival rates of prostheses in the older publications with the survival rates in the newer publications, there was a marked reduction of failures, translating into increased survival rates of implant-supported prostheses in the more recent studies. The difference reached statistical significance ($P = .002$ to $.050$) for all types of prostheses analyzed, except for implant-supported FDPs ($P = .087$).

(Table 3). The most pronounced improvement was seen for screw-retained prostheses, with a 5-year survival rate of 77.6% in the older studies compared with a 5-year survival rate of 96.8% in the newer studies (Table 3).

Esthetic Complications

For implant-supported SCs, there were 12 studies reporting on the esthetic outcome of the treatment. In the older studies the annual rate of implant-supported SCs with semioptimal or unacceptable esthetic outcomes was 3.47%, translating into a 5-year complication rate of 15.9% (Table 4). In the newer studies, the annual rate of esthetic complications was reduced to 1.12%, translating into a 5-year complication rate of 5.4%. The difference in the incidence of esthetic complications between the older and the newer studies did not, however, reach statistical significance ($P = .085$) (Table 4).

Biologic Complications

Peri-implant mucosal lesions were reported in various ways by the different authors. Several studies provided information on soft tissue complications, peri-implantitis, and marginal bone loss, while other studies reported signs of inflammation (pain, redness, swelling, and bleeding) or soft tissue complications, defined as fistula, gingivitis, or hyperplasia.

For implant-supported SCs and implant-supported FDPs, information on biologic complications was extracted from the included publications.

For implant-supported SCs, the annual rate of biologic complications was reduced from 2.56% in the older studies to 1.31% in the new studies, translating into a reduction in 5-year complication rate from 12.0% to 6.4% (Table 3). This difference, however, did not reach statistical significance ($P = .252$). No formal comparison could be made regarding marginal bone levels around implant-supported SCs, because this complication was not reported in the older studies.

For implant-supported FDPs, the incidence of biologic complications increased slightly in the newer studies compared with the older ones. On the other hand, the incidence of substantial bone loss ≥ 2 mm decreased slightly. The annual rate of biologic complications by implant-supported FDPs increased from 1.54% to 1.97%, hence the 5-year complication rate increased from 7.4% to 9.4%. The annual rate of marginal bone loss decreased from 0.68% to




Fig 1 Annual failure rates—overall results. Dots indicate annual failure rates; lines indicate 95% confidence interval.




Fig 2 Annual failure rates of implant-supported SCs.




Fig 3 Annual failure rates of implant-supported FPDs.

0.52%, translating into a 5-year complication rate of 3.3% in the older studies compared with 2.5% in the newer studies. The differences regarding biologic complications between the older and the newer studies for implant-supported FPDs at 5 years did not reach statistical significance ($P = .540, .543$) (Table 4).

Technical Complications

Abutment or Screw Loosening. The annual rate of abutment or screw loosening in the older publications ranged from 0.79% to 6.08%, translating into a 5-year complication rate of 3.9% to 26.2% (Table 5).

The highest incidence of abutment or screw loosening in the older studies was reported for screw-retained prostheses (26.2%) and implant-supported SCs (24.4%), and the lowest complication rate was reported for cemented prostheses (3.9%). In the newer studies, the annual rate of abutment or screw loosening ranged from 0.62% to 2.29%, translating into a 5-year complication rate ranging from 3.1% to 10.8% (Table 5). The highest incidence of abutment or screw loosening in the newer studies was still seen for screw-retained prostheses and the lowest for cemented prostheses. For all types of prostheses, lower incidences of abutment and screw loosening were reported in the newer studies. For screw-retained prostheses and implant-supported SCs, this difference reached statistical significance ($P = .002, .045$) (Table 5).

Abutment or Screw Fractures. The annual rate of abutment or screw fractures in the older publications ranged from 0.16% to 0.44%, translating into a 5-year complication rate of 0.8% to 2.2% (Table 6). In the older studies, this information was not available for screw-retained prostheses and implant-supported FCDs. In the more recent studies, the annual rate of abutment or screw fractures ranged from 0% to 1.20%, translating into a 5-year complication rate between 0% and 5.8% (Table 6). Comparing the overall results in the older and the newer studies in respect to abutment or screw loosening, there was an increase in annual failure rates from 0.27% to 0.56%, representing a change for the 5-year complication rate from 1.3% to 2.8% ($P = .371$). It must, however, be kept in mind that among the older studies, no studies on screw-retained prostheses and implant-supported FCDs were available. When the different types of prostheses were analyzed separately, they all showed a decreased rate of abutment or screw loosening when comparing the older studies with the more recent ones. The difference between the older and the newer studies, however, only reached statistical significance for implant-supported SCs ($P = .029$). The highest 5-year rate of abutment or

screw fractures of 5.8% was reported for implant-supported FCDs. For implant-supported screw-retained prostheses, the 5-year complication rate was 4.1%, compared with a complication rate of 0% for implant-supported cemented prosthesis (Table 6).

Fracture of the Veneering Material.

The annual rate of fracture of the veneering material in the older publications ranged from 0.28% to 4.28%, translating into a 5-year complication rate of 1.4% to 19.2% (Table 7). The highest 5-year rate of fracture of the veneering material in the older studies was reported for implant-supported FDPs. In the newer studies, the annual rate of fracture of the veneering material ranged from 0.64% to 5.82%, translating into a 5-year complication rate ranging between 3.2% and 25.5% (Table 7). The lowest 5-year rate of fracture of the veneering material was reported for implant-supported SCs, and the highest rate was reported for implant-supported FCDs. Comparing the older studies with the newer studies, there was a significant increase in the incidence of fracture of the veneering material for the overall results ($P < .0001$), for the cemented prostheses ($P = .004$), and for the screw-retained prostheses ($P < .0001$). It must, however, be kept in mind that among the older studies, there were no studies reporting on implant-supported FCDs that showed the highest incidence of complications in the newer studies. On the other hand, there was a significant decrease in fracture of the veneering material reported for implant-supported SCs ($P = .054$) and for the implant-supported FDPs ($P = .013$) (Table 7).

Implant Fractures. Implant fractures are a rare complication. For implant-supported SCs, the annual rate of implant fractures was reduced from 0.06% in the older studies to 0.02% in the newer studies, translating into a reduction in the 5-year complication rate from 0.3% to 0.08% (Table 8). This difference did not reach statistical significance ($P = .271$). For implant-supported FDPs the 5-year rate of implant fractures was the same or 0.5% both on the older and the newer studies (Table 8).




Fig 4 Annual failure rates of implant-supported FCDs.




Fig 5 Annual failure rates of implant-supported cemented reconstructions.

Table 4 Comparison of Esthetic and Biologic Complications in Articles Published Through and After 2000

	Published through 2000		Published after 2000		P value
	Annual complication rate (95% CI)	5-year complication rate (95% CI)	Annual complication rate (95% CI)	5-year complication rate (95% CI)	
Esthetic complications, SCs	3.47% (0.86–14.08)	15.9% (4.2–50.5)	1.12% (0.68–1.84)	5.4% (3.3–8.8)	.085
Biologic complications, SCs	2.56% (0.82–7.99)	12.0% (4.0–32.9)	1.31% (0.85–2.05)	6.4% (4.2–9.7)	.252
Biologic complications, FPDs	1.54% (0.74–3.20)	7.4% (3.6–14.8)	1.97% (1.31–2.97)	9.4% (6.3–13.8)	.540
Marginal bone loss \geq 2 mm, SCs	NR	NR	1.31% (0.66–2.58)	6.3% (3.3–12.1)	NA
Marginal bone loss \geq 2 mm, FPDs	0.68% (0.26–1.80)	3.3% (1.3–8.6)	0.52% (0.38–0.70)	2.5% (1.9–3.5)	.543

NR = not reported; NA = not applicable.

Table 5 Comparison of Abutment or Screw Loosening in Articles Published Through and After 2000

	Published through 2000		Published after 2000		P value
	Annual complication rate (95% CI)	5-year complication rate (95% CI)	Annual complication rate (95% CI)	5-year complication rate (95% CI)	
Overall results	1.65% (0.75–3.65)	7.9% (3.7–16.7)	1.82% (1.22–2.73)	8.7% (5.9–12.8)	.826
Cemented reconstructions	0.79% (0.57–1.09)	3.9% (2.8–5.3)	0.62% (0.33–1.21)	3.1% (1.6–5.8)	.530
Screw-retained reconstructions	6.08% (3.79–9.74)	26.2% (17.3–38.6)	2.29% (1.47–3.56)	10.8% (7.1–16.3)	.002
Implant-supported FCDs	NR	NR	1.88% (0.63–5.61)	9.0% (3.1–24.4)	NA
Implant-supported SCs	5.58% (1.19–26.11)	24.4% (5.8–72.9)	1.16% (0.66–2.03)	5.6% (3.2–9.6)	.045
Implant-supported FPDs	1.08% (0.78–1.48)	5.2% (3.8–7.1)	0.81% (0.45–1.46)	4.0% (2.2–7.0)	.387

NR = not reported; NA = not applicable.

Table 6 Comparison of Abutment or Screw Fracture in Articles Published Through and After 2000

	Published through 2000		Published after 2000		P value
	Annual complication rate (95% CI)	5-year complication rate (95% CI)	Annual complication rate (95% CI)	5-year complication rate (95% CI)	
Overall results	0.27% (0.05–1.35)	1.3% (0.3–6.5)	0.56% (0.29–1.11)	2.8% (1.4–5.4)	.371
Cemented reconstructions	0.28% (0.05–1.50)	1.4% (0.3–7.2)	0%	0%	NA
Screw-retained reconstructions	NR	NR	0.84% (0.42–1.67)	4.1% (2.1–8.0)	NA
Implant-supported FCDs	NR	NR	1.20% (0.31–4.68)	5.8% (1.5–20.9)	NA
Implant-supported SCs	0.16% (0.05–0.49)	0.8% (0.3–2.4)	0.07% (0.02–0.20)	0.3% (0.1–1.0)	.238
Implant-supported FPDs	0.44% (0.23–0.84)	2.2% (1.1–4.1)	0.16% (0.08–0.32)	0.8% (0.4–1.6)	.029

NR = not reported; NA = not applicable.

DISCUSSION

The aim of this systematic review was to investigate the survival and complication rates of implant-supported prostheses in older studies and compare them with survival and complication rates reported in more recent publications.

With the exception of implant-supported FPDs, implant survival rate was similar in the older and in the

more recent studies. The overall 5-year implant survival rate and the survival rate for implant-supported SCs was high, ranging between 97.1 and 98.6% in both the older and the newer studies. For implant-supported FPDs, the 5-year survival rate even increased over time.

Considering this, what does it mean for daily clinical practice when the survival rate is increased from 93.8% to 96.1%? A survival rate of 93.8% indicates that 1 implant out of 16 was lost, and 96.1% means that

Table 7 Comparison of Fractures of Veneering Material in Articles Published Through and After 2000

	Published through 2000		Published after 2000		P value
	Annual complication rate (95% CI)	5-year complication rate (95% CI)	Annual complication rate (95% CI)	5-year complication rate (95% CI)	
Overall results	0.31% (0.09–1.03)	1.5% (0.5–5.0)	3.65% (2.55–5.23)	16.7% (12.0–23.0)	.0001
Cemented reconstructions	0.28% (0.08–1.01)	1.4% (0.4–4.9)	2.84% (1.04–7.81)	13.3% (5.0–32.3)	.004
Screw-retained reconstructions	0.76% (0.37–1.55)	3.7% (1.8–7.4)	3.95% (2.74–5.69)	17.9% (12.8–24.8)	.0001
Implant-supported FCDs	NR	NR	5.82% (3.77–9.00)	25.3% (17.2–36.2)	NA
Implant-supported SCs	1.27% (0.67–2.41)	6.2% (3.3–11.4)	0.64% (0.43–0.96)	3.2% (2.1–4.7)	.054
Implant-supported FDPs	4.28% (3.10–5.90)	19.2% (14.4–25.5)	1.60% (0.77–3.30)	7.7% (3.8–15.2)	.013

NR = not reported; NA = not applicable.

Table 8 Comparison of Implant Fractures, Framework Material Fractures, and Loss of Retention in Articles Published Through and After 2000

	Published through 2000		Published after 2000		P value
	Annual complication rate (95% CI)	5-year complication rate (95% CI)	Annual complication rate (95% CI)	5-year complication rate (95% CI)	
Implant fractures, SCs	0.06% (0.008–0.52)	0.3% (0.04–2.6)	0.02% (0.004–0.07)	0.08% (0.02–0.35)	.271
Implant fractures, FDPs	0.10% (0.03–0.30)	0.5% (0.2–1.5)	0.11% (0.4–0.30)	0.5% (0.2–1.5)	.912
Fractures of the framework, FDPs	0.19% (0.08–0.50)	1.0% (0.4–2.4)	0.04% (0.005–0.27)	0.2% (0.02–1.4)	.128
Loss of retention, SCs	1.52% (0.56–4.17)	7.3% (2.7–18.8)	0.63% (0.30–1.29)	3.1% (1.5–6.3)	.118

NR = not reported; NA = not applicable.

Table 9 Comparison of Combined Complication Rates in Articles Published Through and After 2000

	Published through 2000		Published after 2000		P value
	Annual complication rate (95% CI)	5-year complication rate (95% CI)	Annual complication rate (95% CI)	5-year complication rate (95% CI)	
Overall results	3.21% (1.85–5.56)	14.8% (8.8–24.3)	6.33% (4.78–8.38)	27.1% (21.3–34.2)	.028
Cemented reconstructions	2.32% (1.50–3.57)	10.9% (7.2–16.3)	3.55% (2.05–6.17)	16.3% (9.7–26.5)	.225
Screw-retained reconstructions	8.10% (4.02–16.32)	33.3% (18.2–55.8)	7.42% (5.5–10.1)	31.0% (23.9–39.7)	.808
Implant-supported FCDs	NR	NR	15.19% (9.62–24.0)	53.2% (38.2–69.9)	NA
Implant-supported FDPs	10.46% (10.20–10.73)	40.1% (39.9–41.5)	6.37% (4.41–9.20)	27.3% (19.8–36.9)	.005

NR = not reported; NA = not applicable.

1 implant out of 26 was lost. To simplify, for example, 99% survival means loss of 1 implant out of 100, and 90% means loss of 1 implant out of 10. Hence, it has a major influence on the daily practice whether the survival and/or the success rate of an implant-supported prosthesis is 90% or 99%.

For all groups of implant-supported prostheses, there was a substantial to significant improvement in survival rates comparing the older studies with the newer studies. In the older studies, the 5-year survival rates were between 77.6% and 95.2%, compared with

survival rates between 95.8% and 97.9% in the newer studies. The most significant improvement was reported for screw-retained implant-supported prostheses.

A positive improvement was also seen regarding esthetic outcomes comparing the older to the newer studies. This might represent a positive learning curve regarding improved understanding of biologic principles that must be respected during implant treatment in areas of esthetic priority. This might also represent a positive influence of new materials like ceramics, most specifically zirconia, that make it possible to improve

the esthetic outcome of the treatment.^{1,103} The results regarding biologic complications were not consistent. For implant-supported SCs, the incidence of biologic complications decreased from the older studies compared with the newer studies. For implant-supported FDPs, there was a slight increase in biologic complications and a slight decrease in the number of implants with substantial marginal bone loss. The changes in esthetic outcomes and biologic complications did not reach statistical significance.

For most of the implant-supported prostheses, there were slightly to significantly fewer incidences of screw or abutment loosening and fractures, again displaying an improvement of the materials and methods. For screw-retained prostheses, the rate of screw or abutment loosening was reduced from 26.2% to 10.8%, and for implant-supported SCs the complication rate was reduced from 24.4% to 5.6%. One of the reasons for the significant reduction in screw or abutment loosening was a clear outlier among the older studies, Henry et al,¹⁰⁴ reporting on the first generation of single crowns on Brånemark implants. This group reported on titanium screws replaced with new gold abutment screws and new abutments replaced with older ones, resulting in dramatically reduced screw loosening.

Fracture of the veneering material was the most frequently reported technical complication. Comparing the rate of this complication within the older and the newer studies, the results varied significantly. For implant-supported SCs and implant-supported FDPs with at least 5 years of follow-up time, there was a significant decrease in the incidence of veneering material fractures. One of the reasons for this is probably that several studies reporting on implant-supported prostheses with gold-framework and acrylic veneers are included in the group of older studies. It has been demonstrated in previous systematic reviews^{4,14} that implant-supported prostheses with acrylic veneers have a significantly lower survival rate than implant-supported metal-ceramic prostheses. On the other hand, the fracture rate of veneering material reported in studies with shorter follow-up time was significantly increased in the newer studies compared with the older studies. The risk of fracture of the veneering material was increased with the size of the reconstruction. The lowest 5-year complication rate (3.2%) was reported for implant-supported SCs, and the highest complication rate (25.3%) was reported for implant-supported FCDs. It is difficult to speculate what could be the reason for increased rate of fractures of the veneering material. One explanation could be a tendency to evaluate and report complications in more detail in recent publications. A minor ceramic chipping is a typical complication that could go unnoticed if the clinical examiner is not carefully investigating the prostheses. This could

also explain the fact that the total number of technical complications was significantly higher in the newer studies compared with the older studies. Another explanation could be the increased application of more delicate types of prostheses, eg, zirconia- or titanium-based implant FDPs. The veneering ceramics for these types of framework materials exhibited high rates of chipping in clinical studies.

The high rate of technical complications must be given serious consideration. The 5-year rate of technical complications ranged from 16.3% to 53.2%. The lowest rate was reported for cemented prostheses and the highest rate was reported for implant-supported FCDs, where every second prosthesis had a technical complication of some kind. Since the latter observation is only based on very few studies, it has to be interpreted with caution. Specific clinician- or technician-based factors might be one possible reason for these complications.

The 41 meta-analyses performed are based on 139 clinical studies reporting on 8,193 implant-supported prostheses. Therefore, it can be concluded that the results are based on substantial material size. Another strength of the present systematic review is that the methodology used is well standardized in the way the search strategy was performed, the data extraction, and how the statistical approach was performed. Due to the fact that there was a substantial heterogeneity among the included studies, it was decided to use the robust Poisson regression, which incorporated heterogeneity among the studies.

One limitation of this review is that it was mainly based on studies that were conducted in an institutional environment, such as university or specialized implant clinics. Therefore, the long-term outcomes observed cannot be generalized to dental services provided in private practice. A further limitation is that the published information did not allow estimating annual failure rates separately for different time periods or years after insertion of the prosthesis. Thus, it was not possible to estimate whether annual failure rates increased over time. One of the limitations of the present systematic review was that both prospective and retrospective cohort studies and case series were included. To assess the influence of study design, the results from prospective and retrospective studies have been analyzed separately in a recent systematic review.^{5,14} In two of the analyses, no influence of study effect could be seen, but in the third analysis higher survival rates were reported for retrospective studies. Hence, it was difficult to draw any robust conclusions regarding the influence of including retrospective studies in the analysis. In the present systematic review, the study design should not be a problem as long as the distribution of retrospective and prospective studies is similar between the older and the more recent study groups.

Table 10 Comparison of Failure and Complication Rates of Implant-Supported SCs and FDPs in Articles Published Before 2000, from 2000 to 2005, and After 2005

	Published before 2000		Published 2000–2005	
	Annual failure/ complication rate (95% CI)	5-year survival/ complication rate (95% CI)	Annual failure/ complication rate (95% CI)	5-year survival/ complication rate (95% CI)
Implant survival, SCs	0.52% (0.35–0.79)	97.4% (96.1–98.3)	0.61% (0.36–1.04)	97.0% (95.0–98.2)
Implant survival, FDPs	1.35% (1.12–1.62)	93.5% (92.2–94.5)	0.81% (0.53–1.22)	96.0% (94.1–97.4)
Survival of implant-supported SCs	1.54% (0.97–2.44)	92.6% (88.5–95.3)	0.83% (0.53–1.30)	95.9% (93.7–97.4)
Survival of implant-supported FDPs	1.57% (0.80–3.07)	92.4% (85.8–96.1)	0.63% (0.49–0.83)	96.9% (95.9–97.6)
Biologic complications, SCs	2.69% (0.85–8.57)	12.6% (4.1–34.8)	1.44% (0.83–2.51)	6.9% (4.0–11.8)
Esthetic complications, SCs	3.47% (0.86–14.08)	15.9% (4.2–50.5)	1.17% (0.40–3.41)	5.7% (2.0–15.7)
Implant fractures, SCs	0.08% (0.009–0.72)	0.4% (0.04–3.54)	0.03% (0.004–0.2)	0.1% (0.02–0.99)
Implant fractures, FDPs	1.35% (1.12–1.62)	6.5% (5.5–7.8)	0.81% (0.53–1.22)	4.0% (2.6–5.9)
Abutment or screw loosening, SCs	6.81% (1.43–32.26)	28.8% (6.9–80.1)	0.95% (0.37–2.49)	4.7% (1.8–11.7)
Fractures of the veneering material, SCs	1.27% (0.67–2.41)	6.2% (3.3–11.4)	0.49% (0.24–1.01)	2.4% (1.2–4.9)
Fractures of the veneering material, FDPs	4.66% (3.54–6.14)	20.8% (16.2–26.4)	1.29% (0.68–2.43)	6.3% (3.4–11.5)
Total number of complications, FDPs	10.46% (10.2–10.73)	40.1% (39.9–41.5)	8.10% (3.50–18.74)	33.3% (16.1–60.8)

*Published before 2000 vs published 2000–2005.

†Published before 2000 vs published after 2005.

CONCLUSIONS

Despite of high survival rate of implant-supported prostheses and substantial improvements within implant dentistry over time, esthetic, biologic, and technical complications are still frequent. This, in turn, means that a substantial amount of chair time has to be accepted by the patient and dental services. The present systematic review demonstrated in many aspects a positive learning curve in implant dentistry, represented by lower failure and complication rates reported in more recent clinical studies.

It is, however, of outmost importance that the industry, the scientific community, and clinicians worldwide work together to identify failures, complications, and weaknesses in implant dentistry and develop solutions that make implant treatment an even more predictable and safe therapeutic option.

ACKNOWLEDGMENTS

The authors reported no conflicts of interest related to this study.

REFERENCES

- Jung RE, Pjetursson BE, Glauser R, Zembic A, Zwahlen M, Lang NP. A systematic review of the 5-year survival and complication rates of implant-supported single crowns. *Clin Oral Implants Res* 2008;19: 119–130.
- Jung RE, Zembic A, Pjetursson BE, Zwahlen M, Thoma DS. Systematic review of the survival rate and the incidence of biological, technical, and aesthetic complications of single crowns on implants reported in longitudinal studies with a mean follow-up of 5 years. *Clin Oral Implants Res* 2012;23(suppl 6):2–21.
- Pjetursson BE, Tan K, Lang NP, Bragger U, Egger M, Zwahlen M. A systematic review of the survival and complication rates of fixed partial dentures (FPDs) after an observation period of at least 5 years. *Clin Oral Implants Res* 2004;15:625–642.
- Pjetursson BE, Bragger U, Lang NP, Zwahlen M. Comparison of survival and complication rates of tooth-supported fixed dental prostheses (FDPs) and implant-supported FDPs and single crowns (SCs). *Clin Oral Implants Res* 2007;18(suppl 3):97–113.
- Pjetursson BE, Thoma D, Jung R, Zwahlen M, Zembic A. A systematic review of the survival and complication rates of implant-supported fixed dental prostheses (FDPs) after a mean observation period of at least 5 years. *Clin Oral Implants Res* 2012;23(suppl 6):22–38.
- Papaspyridakos P, Chen CJ, Singh M, Weber HP, Gallucci GO. Success criteria in implant dentistry: A systematic review. *J Dent Res* 2012;91: 242–248.
- Klinge B, Meyle J. Peri-implant tissue destruction. The Third EAO Consensus Conference 2012. *Clin Oral Implants Res* 2012;23(suppl 6): 108–110.
- Salvi GE, Bragger U. Mechanical and technical risks in implant therapy. *Int J Oral Maxillofac Implants* 2009;24(suppl):69–85.

Published after 2005

Annual failure/ complication rate 95% CI)	5-year survival/ complication rate (95% CI)	P value
0.55% (0.36–0.84)	97.3% (95.9–98.2)	.636*, .839†
0.78% (0.51–1.20)	96.2% (94.2–97.5)	.021, .007
0.53% (0.28–1.03)	97.4% (95.0–98.6)	.040, .006
0.83% (0.40–1.70)	95.9% (91.9–98.0)	.008, .166
1.23% (0.66–2.28)	5.9% (3.2–10.8)	.289, .196
1.09% (0.61–1.94)	5.3% (3.0–9.3)	.166, .083
0.01% (0.001–0.09)	0.05% (0.006–0.5)	.467, .172
0.78% (0.51–1.20)	3.8% (2.5–5.8)	.021, .007
1.19% (0.61–2.35)	5.8% (3.0–11.1)	.020, .026
0.68% (0.44–1.05)	3.3% (2.2–5.1)	.025, .084
2.08% (0.59–7.32)	9.9% (2.9–30.7)	.0001, .170
5.85% (3.90–8.78)	25.4% (17.7–35.5)	.441, .0001

9. Chaar MS, Att W, Strub JR. Prosthetic outcome of cement-retained implant-supported fixed dental restorations: A systematic review. *J Oral Rehabil* 2011;38:697–711.
10. Sailer I, Muhlemann S, Zwahlen M, Hammerle CH, Schneider D. Cemented and screw-retained implant reconstructions: A systematic review of the survival and complication rates. *Clin Oral Implants Res* 2012;23(suppl 6):163–201.
11. Larsson C, Vult von Steyern P. Five-year follow-up of implant-supported Y-TZP and ZTA fixed dental prostheses. A randomized, prospective clinical trial comparing two different material systems. *Int J Prosthodont* 2010;23:555–561.
12. Sailer I, Gottner B, Kanelb S, Hammerle CH. Randomized controlled clinical trial of zirconia-ceramic and metal-ceramic posterior fixed dental prostheses: A 3-year follow-up. *Int J Prosthodont* 2009;22:553–560.
13. Sax C, Hammerle CH, Sailer I. 10-year clinical outcomes of fixed dental prostheses with zirconia frameworks. *Int J Comput Dent* 2011;14:183–202.
14. Pjetursson BE, Tan K, Lang NP, Bragger U, Egger M, Zwahlen M. A systematic review of the survival and complication rates of fixed partial dentures (FPDs) after an observation period of at least 5 years. *Clin Oral Implants Res* 2004;15:667–676.
15. Kirkwood BR, Sterne JAC. Poisson regression. *Essential Medical Statistics* 2003:249–262.
16. Kirkwood BR, Sterne JAC. Survival analysis: Displaying and comparing survival patterns. *Essential Medical Statistics* 2003:272–286.
17. Pieri F, Aldini NN, Fini M, Marchetti C, Corinaldesi G. Preliminary 2-year report on treatment outcomes for 6-mm-long implants in posterior atrophic mandibles. *Int J Prosthodont* 2012;25:279–289.
18. Gallucci GO, Grutter L, Nedir R, Bischof M, Belser UC. Esthetic outcomes with porcelain-fused-to-ceramic and all-ceramic single-implant crowns: A randomized clinical trial. *Clin Oral Implants Res* 2011;22:62–69.
19. Di Giacomo GA, da Silva JV, da Silva AM, Paschoal GH, Cury PR, Szarf G. Accuracy and complications of computer-designed selective laser sintering surgical guides for flapless dental implant placement and immediate definitive prosthesis installation. *J Periodontol* 2012;83:410–419.
20. Hjalmarsson L, Smedberg JI, Pettersson M, Jemt T. Implant-level prostheses in the edentulous maxilla: A comparison with conventional abutment-level prostheses after 5 years of use. *Int J Prosthodont* 2011;24:158–167.
21. Schneider D, Grunder U, Ender A, Hammerle CH, Jung RE. Volume gain and stability of peri-implant tissue following bone and soft tissue augmentation: 1-year results from a prospective cohort study. *Clin Oral Implants Res* 2011;22:28–37.
22. Crespi R, Cappare P, Gherlone E. Osteotome sinus floor elevation and simultaneous implant placement in grafted biomaterial sockets: 3 years of follow-up. *J Periodontol* 2010;81:344–349.
23. Eliasson A, Narby B, Ekstrand K, Hirsch J, Johansson A, Wennerberg A. A 5-year prospective clinical study of submerged and nonsubmerged Paragon system implants in the edentulous mandible. *Int J Prosthodont* 2010;23:231–238.
24. Larsson C, von Steyern PV (2010). Five-year follow-up of implant-supported Y-TZP and ZTA fixed dental prostheses. A randomized, prospective clinical trial comparing two different material systems. *Int J Prosthodont* 2010;23:555–561.
25. Schwarz S, Gabbert O, Hassel AJ, Schmitter M, Seche C, Rammelsberg P. Early loading of implants with fixed dental prostheses in edentulous mandibles: 4.5-year clinical results from a prospective study. *Clin Oral Implants Res* 2010;21:284–289.
26. Gallucci GO, Doughtie CB, Hwang JW, Fiorellini JP, Weber HP. Five-year results of fixed implant-supported rehabilitations with distal cantilevers for the edentulous mandible. *Clin Oral Implants Res* 2009;20:601–607.
27. Jemt T. Cemented CeraOne and porcelain fused to TiAdapt abutment single-implant crown restorations: A 10-year comparative follow-up study. *Clin Implant Dent Relat Res* 2009;11:303–310.
28. Ortorp A, Jemt T. CNC-milled titanium frameworks supported by implants in the edentulous jaw: A 10-year comparative clinical study. *Clin Implant Dent Relat Res* 2009;14:88–99.
29. Ortorp A, Jemt T. Early laser-welded titanium frameworks supported by implants in the edentulous mandible: A 15-year comparative follow-up study. *Clin Implant Dent Relat Res* 2009;11:311–322.
30. Zembic A, Sailer I, Jung RE, Hammerle CH. Randomized-controlled clinical trial of customized zirconia and titanium implant abutments for single-tooth implants in canine and posterior regions: 3-year results. *Clin Oral Implants Res* 2009;20:802–808.
31. Akca K, Cehreli MC. Two-year prospective follow-up of implant/tooth-supported versus frestanding implant-supported fixed partial dentures. *Int J Periodontics Restorative Dent* 2008;28:593–599.
32. Cannizzaro G, Leone M, Consolo U, Ferri V, Esposito M. Immediate functional loading of implants placed with flapless surgery versus conventional implants in partially edentulous patients: A 3-year randomized controlled clinical trial. *Int J Oral Maxillofac Implants* 2008;23:867–875.
33. Cecchinato D, Bengazi F, Blasi G, Botticelli D, Cardarelli I, Gualini F. Bone level alterations at implants placed in the posterior segments of the dentition: Outcome of submerged/non-submerged healing. A 5-year multicenter, randomized, controlled clinical trial. *Clin Oral Implants Res* 2008;19:429–431.
34. Fischer K, Stenberg T, Hedin M, Sennerby L. Five-year results from a randomized, controlled trial on early and delayed loading of implants supporting full-arch prosthesis in the edentulous maxilla. *Clin Oral Implants Res* 2008;19:433–441.
35. Halg GA, Schmid J, Hammerle CH. Bone level changes at implants supporting crowns or fixed partial dentures with or without cantilevers. *Clin Oral Implants Res* 2008;19:983–990.
36. Ortorp A, Jemt T. Laser-welded titanium frameworks supported by implants in the partially edentulous mandible: A 10-year comparative follow-up study. *Clin Implant Dent Relat Res* 2008;10:128–139.

37. Schropp L, Isidor F. Clinical outcome and patient satisfaction following full-flap elevation for early and delayed placement of single-tooth implants: A 5-year randomized study. *Int J Oral Maxillofac Implants* 2008;23:733–743.
38. Cooper LF, Ellner S, Moriarty J, et al. Three-year evaluation of single-tooth implants restored 3 weeks after 1-stage surgery. *Int J Oral Maxillofac Implants* 2007;22:791–800.
39. Crespi R, Cappare P, Gherlone E, Romanos GE. Immediate occlusal loading of implants placed in fresh sockets after tooth extraction. *Int J Oral Maxillofac Implants* 2007;22:955–962.
40. Nordin T, Graf J, Frykholm A, Hellden L. Early functional loading of sand-blasted and acid-etched (SLA) Straumann implants following immediate placement in maxillary extraction sockets. Clinical and radiographic result. *Clin Oral Implants Res* 2007;18:441–451.
41. Drago CJ, Lazzara RJ. Immediate occlusal loading of Osseotite implants in mandibular edentulous patients: A prospective observational report with 18-month data. *J Prosthodont* 2006;15:187–194.
43. Turkyilmaz I. A 3-year prospective clinical and radiologic analysis of early loaded maxillary dental implants supporting single-tooth crowns. *Int J Prosthodont* 2006;19:389–390.
42. Eliasson A, Eriksson T, Johansson A, Wennerberg A. Fixed partial prostheses supported by 2 or 3 implants: A retrospective study up to 18 years. *Int J Oral Maxillofac Implants* 2006;21:567–574.
44. Abboud M, Koeck B, Stark H, Wahl G, Paillon R. Immediate loading of single-tooth implants in the posterior region. *Int J Oral Maxillofac Implants* 2005;20:61–68.
45. Bragger U, Karoussis I, Persson R, Pjetursson B, Salvi G, Lang N. Technical and biological complications/failures with single crowns and fixed partial dentures on implants: A 10-year prospective cohort study. *Clin Oral Implants Res* 2005;16:326–334.
46. Rasmusson L, Roos J, Bystedt H. A 10-year follow-up study of titanium dioxide-blasted implants. *Clin Implant Dent Relat Res* 2005;7:36–42.
47. van Steenberghe D, Glauser R, Blomback U, et al. A computed tomographic scan-derived customized surgical template and fixed prosthesis for flapless surgery and immediate loading of implants in fully edentulous maxillae: A prospective multicenter study. *Clin Implant Dent Relat Res* 2005;7(suppl 1):s111–s120.
48. Astrand P, Engquist B, Anzen B, et al. A three-year follow-up report of a comparative study of ITI dental implants and Branemark System implants in the treatment of the partially edentulous maxilla. *Clin Implant Dent Relat Res* 2004;6:130–141.
49. Astrand P, Engquist B, Dahlgren S, Grondahl K, Engquist E, Feldmann H. Astra Tech and Branemark system implants: A 5-year prospective study of marginal bone reactions. *Clin Oral Implants Res* 2004;15:413–420.
50. Glauser R, Sailer I, Wohlwend A, Studer S, Schibli M, Scharer P. Experimental zirconia abutments for implant-supported single-tooth restorations in esthetically demanding regions: 4-year results of a prospective clinical study. *Int J Prosthodont* 2004;17:285–290.
51. Gotfredsen K. A 5-year prospective study of single-tooth replacements supported by the Astra Tech implant: A pilot study. *Clin Implant Dent Relat Res* 2004;6:1–8.
52. Henriksson K, Jemt T. Measurements of soft tissue volume in association with single-implant restorations: A 1-year comparative study after abutment connection surgery. *Clin Implant Dent Relat Res* 2004;6:181–189.
53. Drago CJ. A clinical study of the efficacy of gold-tite square abutment screws in cement-retained implant restorations. *Int J Oral Maxillofac Implants* 2003;18:273–278.
54. Duncan JP, Nazarova E, Vogiatzi T, Taylor TD. Prosthodontic complications in a prospective clinical trial of single-stage implants at 36 months. *Int J Oral Maxillofac Implants* 2003;18:561–565.
55. Hellden L, Ericson G, Elliot A, et al. A prospective 5-year multicenter study of the Cresco implantology concept. *Int J Prosthodont* 2003;16:554–562.
56. Andersen E, Saxegaard E, Knutsen BM, Haanaes HR. A prospective clinical study evaluating the safety and effectiveness of narrow-diameter threaded implants in the anterior region of the maxilla. *Int J Oral Maxillofac Implants* 2001;16:217–224.
57. Bambini F, Lo Muzio L, Procaccini M. Retrospective analysis of the influence of abutment structure design on the success of implant unit. A 3-year controlled follow-up study. *Clin Oral Implants Res* 2001;12:319–324.
58. Johnson RH, Persson GR. A 3-year prospective study of a single-tooth implant—Prosthodontic complications. *Int J Prosthodont* 2001;14:183–189.
59. Astrand P, Anzen B, Karlsson U, Sahlholm S, Svardstrom P, Hellem S. Nonsubmerged implants in the treatment of the edentulous upper jaw: A prospective clinical and radiographic study of ITI implants—Results after 1 year. *Clin Implant Dent Relat Res* 2000;2:166–174.
60. Palmer RM, Palmer PJ, Smith BJ. A 5-year prospective study of Astra single tooth implants. *Clin Oral Implants Res* 2000;11:179–182.
61. Romanos GE, Nentwig GH. Single molar replacement with a progressive thread design implant system: A retrospective clinical report. *Int J Oral Maxillofac Implants* 2000;15:831–836.
62. Scholander S. A retrospective evaluation of 259 single-tooth replacements by the use of Branemark implants. *Int J Prosthodont* 1999;12:483–491.
63. Wannfors K, Smedberg JI. A prospective clinical evaluation of different single-tooth restoration designs on osseointegrated implants. A 3-year follow-up of Branemark implants. *Clin Oral Implants Res* 1999;10:453–458.
64. Andersson B, Odman P, Lindvall AM, Brånemark PI. Five-year prospective study of prosthodontic and surgical single-tooth implant treatment in general practices and at a specialist clinic. *Int J Prosthodont* 1998;11:351–355.
65. Andersson B, Odman P, Lindvall AM, Brånemark PI. Cemented single crowns on osseointegrated implants after 5 years: Results from a prospective study on CeraOne. *Int J Prosthodont* 1998;11:212–218.
66. Scheller H, Urgell JP, Kultje C, et al. A 5-year multicenter study on implant-supported single crown restorations. *Int J Oral Maxillofac Implants*. 1998;13:212–218.
67. Karlsson U, Gotfredsen K, Olsson C. Single-tooth replacement by osseointegrated Astra Tech dental implants: A 2-year report. *Int J Prosthodont* 1997;10:318–324.
68. Kempainen P, Eskola S, Ylipaavalniemi P. A comparative prospective clinical study of two single-tooth implants: A preliminary report of 102 implants. *J Prosthet Dent* 1997;77:382–387.
69. Henry PJ, Rosenberg IR, Bills IG, et al. Osseointegrated implants for single tooth replacement in general practice: A 1-year report from a multicentre prospective study. *Aust Dent J* 1995;40:173–181.
70. Jung RE, Zaugg B, Philipp AO, Truninger TC, Siegenthaler DW, Hämmerle CH. A prospective controlled clinical trial evaluating the clinical radiological and aesthetic outcome after 5 years of immediately placed implants in sockets exhibiting pericapital pathology. *Clin Oral Implants Res* 2013;24:839–846.
71. Schneider D, Witt L, Hämmerle CH. Influence of the crown-to-implant length ratio on the clinical performance of implants supporting single crown restorations: A cross-sectional retrospective 5-year investigation. *Clin Oral Implants Res* 2012;23:169–174.
72. Krennmair G, Seemann R, Schmidinger S, Ewers R, Piehslinger E. Clinical outcome of root-shaped dental implants of various diameters: 5-year results. *Int J Oral Maxillofac Implants* 2010;25:357–366.
73. Schmidlin K, Schnell N, Steiner S, et al. Complication and failure rates in patients treated for chronic periodontitis and restored with single crowns on teeth and/or implants. *Clin Oral Implants Res* 2010;21:550–557.
74. Urdaneta RA, Rodriguez S, McNeil DC, Weed M, Chuang SK. The effect of increased crown-to-implant ratio on single-tooth locking-taper implants. *Int J Oral Maxillofac Implants* 2010;25:729–743.
75. Vigolo P, Givani A. Platform-switched restorations on wide-diameter implants: A 5-year clinical prospective study. *Int J Oral Maxillofac Implants* 2009;24:103–109.
76. Degidi M, Iezzi G, Perrotti V, Piattelli A. Comparative analysis of immediate functional loading and immediate nonfunctional loading to traditional healing periods: A 5-year follow-up of 550 dental implants. *Clin Implant Dent Relat Res* 2009;11:257–266.
77. Kreissl ME, Gerdts T, Mucche R, Heydecke G, Strub JR. Technical complications of implant-supported fixed partial dentures in partially edentulous cases after an average observation period of 5 years. *Clin Oral Implants Res* 2007;18:720–726.
78. Wennström JL, Ekestubbe A, Gröndahl K, Karlsson S, Lindhe J. Implant-supported single-tooth restorations: A 5-year prospective study. *J Clin Periodontol* 2005;32:567–574.

79. Bernard JP, Schatz JP, Christou P, Belsler U, Kiliaridis S. Long-term vertical changes of the anterior maxillary teeth adjacent to single implants in young and mature adults. A retrospective study. *J Clin Periodontol* 2004;31:1024–1028.
80. Haas R, Polak C, Fürhauser R, Mailath-Pokorny G, Dörtbudak O, Watzek G. A long-term follow-up of 76 Branemark single-tooth implants. *Clin Oral Implants Res* 2002;13:38–43.
81. Mericske-Stern R, Grütter L, Rösch R, Mericske E. Clinical evaluation and prosthetic complications of single tooth replacements by non-submerged implants. *Clin Oral Implants Res* 2001;12:309–318.
82. Polizzi G, Fabbro S, Furri M, Herrmann I, Squarzone S. Clinical application of narrow Branemark System implants for single-tooth restorations. *Int J Oral Maxillofac Implants*. 1999;14:496–503.
83. Ridell A, Gröndahl K, Sennerby L. Placement of Branemark implants in the maxillary tuber region: Anatomical considerations, surgical technique and long-term results. *Clin Oral Implants Res* 2009;20:94–98
84. Ormianer Z, Palt, A. Long-term clinical evaluation of tapered multi-threaded implants: Results and influences of potential risk factors. *J Oral Implantol*. 2006;32:300–307.
85. Romeo E, Ghisolfi M, Rozza R, Chiapasco M, Lops D. Short (8-mm) dental implants in the rehabilitation of partial and complete edentulism: A 3- to 14-year longitudinal study. *Int J Prosthodont* 2006;19:586–592.
86. Degidi M, Piattelli A. A 7-year follow-up of 93 immediately loaded titanium dental implants. *J Oral Implantol* 2005;31:25–31.
87. Becker CM. Cantilever fixed prostheses utilizing dental implants: A 10-year retrospective analysis. *Quintessence Int* 2004;35:437–441.
88. Wennström JL, Ekstubb A, Gröndahl K, Karlsson S, Lindhe J. Oral rehabilitation with implant-supported fixed partial dentures in periodontitis-susceptible subjects. A 5-year prospective study. *J Clin Periodontol* 2004;31:713–724.
89. Andersson B, Glauser R, Maglione M, Taylor A. Ceramic implant abutments for short-span FDPs: A prospective 5-year multicenter study. *Int J Prosthodont* 2003;16:640–646.
90. Jemt T, Henry P, Lindén B, Naert I, Weber H, Wendelhag I. Implant-supported laser-welded titanium and conventional cast frameworks in the partially edentulous jaw: A 5-year prospective multicenter study. *Int J Prosthodont* 2003;16:415–421.
91. Preiskel HW, Tsolka P. Cement- and screw-retained implant-supported Prostheses: Up to 10 years of follow-up of a new design. *Int J Oral Maxillofac Implants* 2004;19:87–91.
92. Naert I, Koutsikakis G, Duyck J, Quirynen M, Jacobs R, van Steenberghe D. Biologic outcome of implant-supported restorations in the treatment of partial edentulism Part I: A longitudinal clinical evaluation. *Clin Oral Implants Res* 2002;13:381–389; Naert I, Koutsikakis G, Quirynen M, Duyck J, van Steenberghe D, Jacobs R. Biologic outcome of implant-supported restorations in the treatment of partial edentulism Part 2: A longitudinal radiographic evaluation. *Clin Oral Implants Res* 2002;13:390–395.
93. Brägger U, Karoussis I, Persson R, Pjetursson B, Salvi G, Lang N. Technical and biological complications/failures with single crowns and fixed partial dentures on implants: A 10-year prospective cohort study. *Clin Oral Implants Res* 2005;16:326–334.
94. Gotfredsen K, Karlsson U. A prospective 5-year study of fixed partial-prostheses supported by implants with machined and TiO₂-blasted surface. *J Prosthodont* 2001;10:2–7.
95. Mengel R, Schröder T, Flores-de-Jacoby L. Osseointegrated implants in patients treated for generalized chronic periodontitis and generalized aggressive periodontitis: 3- and 5-year results of a prospective long-term study. *J Periodontol* 2001;72:977–989.
96. Behneke A, Behneke N, d'Hoedt B. The longitudinal clinical effectiveness of ITI solid screw implants in partially edentulous patients: A 5 year follow-up report. *Int J Oral Maxillofac Implants* 2000;15:633–645.
97. Hosny M, Duyck J, van Steenberghe D, Naert I. Within-subject comparison between connected and nonconnected tooth-to-implant fixed partial prostheses: Up to 14-year follow-up study. *Int J Prosthodont* 2000;13:340–346.
98. Wennerberg A, Jemt T. Complications in partially edentulous implant patients: A 5-year retrospective follow-up study of 133 patients supplied with unilateral maxillary prostheses. *Clin Implant Dent Relat Res* 1999;1:49–56.
99. Ortop A, Jemt T. Clinical experiences of implant-supported prostheses with laser-welded titanium frameworks in the partially edentulous jaw: A 5-year follow-up study. *Clin Implant Dent Relat Res* 1999;1:84–91.
100. Wyatt CC, Zarb GA. Treatment outcomes of patients with implant-supported fixed partial prostheses. *Int J Oral Maxillofac Implants* 1998;13:204–211.
101. Olsson M, Gunne J, Astrand P, Borg K. Bridges supported by free-standing implants versus bridges supported by tooth and implant A five-year prospective study. *Clin Oral Implants Res* 1995;6:114–121.
102. Lekholm U, van Steenberghe D, Herrmann I, et al. Osseointegrated implants in the treatment of partially edentulous jaws: A prospective 5-year multicenter study. *Int J Oral Maxillofac Implants* 1994;9:627–635.
103. Sailer I, Philipp A, Zembic A, Pjetursson BE, Hammerle CH, Zwahlen M. A systematic review of the performance of ceramic and metal implant abutments supporting fixed implant reconstructions. *Clin Oral Implants Res* 2009;20(suppl 4):4–31.
104. Henry PJ, Laney WR, Jemt T, Harris D, Krogh PH, Polizzi G, et al. Osseointegrated implants for single-tooth replacement: A prospective 5-year multicenter study. *Int J Oral Maxillofac Implants* 1996;11:450–455.
105. Attard N, Zarb GA. Implant prosthodontic management of posterior partial edentulism: Long-term follow-up of a prospective study. *J Can Dent Assoc* 2002;68:118–124.
106. Bianchi AE, Sanfilippo F. Single-tooth replacement by immediate implant and connective tissue graft: A 1–9-year clinical evaluation. *Clin Oral Implants Res* 2004;15:269–277.
107. Bonde MJ, Stokholm R, Isidor F, Schou S. Outcome of implant-supported single-tooth replacements performed by dental students. A 10-year clinical and radiographic retrospective study. *Eur J Oral Implantol* 2010;3:37–46.
108. Bornstein MM, Schmid B, Belsler UC, Lussi A, Buser D. Early loading of non-submerged titanium implants with a sandblasted and acid-etched surface. 5-year results of a prospective study in partially edentulous patients. *Clin Oral Implants Res* 2005;16:631–638.
109. Cho SC, Small PN, Elian N, Tarnow D. Screw loosening for standard and wide diameter implants in partially edentulous cases: 3- to 7-year longitudinal data. *Implant Dent* 2004;13:245–250.
110. De Boever AL, Keersmaekers K, Vanmaele G, Kerschbaum T, Theuniers G, De Boever JA. Prosthetic complications in fixed osseous implant-borne reconstructions after an observations period of at least 40 months. *J Oral Rehabil* 2006;33:833–839.
111. De Leonardis D, Garg AK, Pecora GE. Osseointegration of rough acid-etched titanium implants: 5-year follow-up of 100 minimatic implants. *Int J Oral Maxillofac Implants* 1999;14:384–391.
112. De Boever AL, De Boever JA. Guided bone regeneration around non-submerged implants in narrow alveolar ridges: A prospective long-term clinical study. *Clin Oral Implants Res* 2005;16:549–556
113. De Boever AL, Quirynen M, Coucke W, Theuniers G, De Boever JA. Clinical and radiographic study of implant treatment outcome in periodontally susceptible and non-susceptible patients: A prospective long-term study. *Clin Oral Implants Res* 2009;20:1341–1350.
114. Elkhoury JS, McGlumphy EA, Tatakis DN, Beck FM. Clinical parameters associated with success and failure of single-tooth titanium plasma-sprayed cylindrical implants under stricter criteria: A 5-year retrospective study. *Int J Oral Maxillofac Implants* 2005;20:687–694.
115. Fartash B, Arvidson K. Long-term evaluation of single crystal sapphire implants as abutments in fixed prosthodontics. *Clin Oral Implants Res* 1997;8:58–67.
116. Froberg KK, Lindh C, Ericsson I. Immediate loading of Branemark System Implants: a comparison between TiUnite and turned implants placed in the anterior mandible. *Clin Implant Dent Relat Res* 2008;8:187–197.
117. Fugazzotto PA, Vlassis J, Butler B. ITI implant use in private practice: Clinical results with 5,526 implants followed up to 72+ months in function. *Int J Oral Maxillofac Implants* 2004;19:408–412.
118. Gibbard LL, Zarb G. A 5-year prospective study of implant-supported single-tooth replacements. *J Can Dent Assoc* 2002;68:110–116.
119. Gotfredsen K. A 10-Year Prospective Study of Single Tooth Implants Placed in the Anterior Maxilla. *Clin Implant Dent Relat Res* 2012;14:80–87

120. Guncu MB, Aslan Y, Tumer C, Guncu GN, Uysal S2. In-patient comparison of immediate and conventional loaded implants in mandibular molar sites within 12 months. *Clin Oral Implants Res* 2008;19:335–341.
121. Gunne J, Astrand P, Lindh T, Borg K, Olsson M. Tooth-implant and implant supported fixed partial dentures: A 10-year report. *Int J Prosthodont* 1999;12:216–221.
122. Jemt T, Lekholm U. Single implants and buccal bone grafts in the anterior maxilla: Measurements of buccal crestal contours in a 6-year prospective clinical study. *Clin Implant Dent Relat Res* 2005;7:127–135.
123. Jemt T. Single implants in the anterior maxilla after 15 years of follow-up: Comparison with central implants in the edentulous maxilla. *Int J Prosthodont*. 2008;21:400–408.
124. Jung RE, Fenner N, Hammerle CHF, Zitzmann NU. Long-term outcome of implants placed with guided bone regeneration (GBR) using resorbable and non-resorbable membranes after 12 to 14 years. *Clin Oral Implants Res* 2013;24:1065–1073
125. Krieger O, Matuliene G, Hüslér J, Salvi GE, Pjetursson B, Bragger U. Failures and complications in patients with birth defects restored with fixed dental prostheses and single crowns on teeth and/or implants. *Clin Oral Implants Res* 2009;20:809–816
126. Lekholm U, Grondahl K, Jemt T. Outcome of oral implant treatment in partially edentulous jaws followed 20 years in clinical function. *Clin Implant Dent Relat Res* 2006;8:178–186
127. Lekholm U, Gunne J, Henry P, Higuchi K, Lindén U, Bergström C, van Steenberghe D. Survival of the Brånemark implant in partially edentulous jaws: A 10-year prospective multicenter study. *Int J Oral Maxillofac Implants* 1999;14:639–645.
128. Levin L, Pathael S, Dolev E, Schwartz-Arad D. Aesthetic versus surgical success of single dental implants: 1- to 9-year follow-up. *Pract Proced Aesthet Dent* 2005;17:533–538; quiz 540, 566.
129. Levine RA, Clem DS, 3rd, Wilson TG, Jr., Higginbottom F, Solnit G. Multicenter retrospective analysis of the ITI implant system used for single-tooth replacements: Results of loading for 2 or more years. *Int J Oral Maxillofac Implants* 1999;14:516–520.
130. Murphy WM, Absi EG, Gregory MC, Williams KR. A prospective 5-year study of two cast framework alloys for fixed implant-supported mandibular prostheses. *Int J Prosthodont* 2002;15:133–138.
131. MacDonald K, Pharoah M, Todescan R, Deporter D. Use of sintered porous-surfaced dental implants to restore single teeth in the maxilla: A 7- to 9-year follow-up. *Int J Periodontics Restorative Dent* 2009;29:191–199.
132. Matarasso S, Rasperini G, Iorio Siciliano V, Salvi GE, Lang NP, Aglietta M. A 10-year retrospective analysis of radiographic bone-level changes of implants supporting single-unit crowns in periodontally compromised vs. periodontally healthy patients. *Clin Oral Implants Res* 2010;21:898–903.
133. Naert IE, Duyck JA, Hosny MM, Van Steenberghe D. Freestanding and tooth-implant connected prostheses in the treatment of partially edentulous patients. Part I: An up to 15-years clinical evaluation. *Clin Oral Implants Res* 2001;12:237–244.
134. Pikner SS, Gröndahl K, Jemt T, Friberg B. Marginal bone loss at implants: A retrospective, long-term follow-up of turned Branemark System implants. *Clin Implant Dent Relat Res* 2009;11:11–23.
135. Romeo E, Lops D, Margutti E, Ghisolfi M, Chiapasco M, Vogel G. Long-term survival and success of oral implants in the treatment of full and partial arches: A 7-year prospective study with the ITI dental implant system. *Int J Oral Maxillofac Implants* 2004;19:247–259.
136. Taylor RC, McGlumphy EA, Tatakis DN, Beck FM. Radiographic and clinical evaluation of single-tooth BioloK implants: A 5-year study. *Int J Oral Maxillofac Implants* 2004;19:849–854.
137. Thilander B, Odman J, Jemt T. Single implants in the upper incisor region and their relationship to the adjacent teeth. An 8-year follow-up study. *Clin Oral Implants Res* 1999;10:346–355.
138. Vigolo P, Givani A. Clinical evaluation of single-tooth mini-implant restorations: A five-year retrospective study. *J Prosthet Dent* 2000;84:50–54.
139. Wagenberg B, Froum SJ. A retrospective study of 1925 consecutively placed immediate implants from 1988 to 2004. *Int J Oral Maxillofac Implants* 2006;21:71–80.
140. Zafropoulos GG, Deli G, Bartee BK, Hoffmann O. Single-tooth implant placement and loading in fresh and regenerated extraction sockets. Five-year results: A case series using two different implant designs. *J Periodontol* 2010;81:604–615.
141. Zarb JP, Zarb GA. Implant prosthodontic management of anterior partial edentulism: Long-term follow-up of a prospective study. *J Can Dent Assoc* 2002;68:92–96.